



COMPASSION PEDIATRICS

Bedwetting

Did you know that there are about 5 million children in the United States who wet the bed? If your child wets the bed, he or she is not alone.

Though most children are toilet trained between 2 and 4 years of age, some children may not be able to stay dry at night until they are older. Children develop at their own rate. For example, 20% of 5-year-olds, 10% of 7-year-olds, and 5% of 10-year-olds may still wet the bed.

Bedwetting is not a serious medical condition, but it can be a challenging problem for children and parents. Read on to find out more from the American Academy of Pediatrics about bedwetting and what can be done about it.

Causes of Bedwetting

Although not all of the causes of bedwetting are fully understood, the following are some that are possible:

- Your child is a deep sleeper and does not awaken to the signal of a full bladder.
- Your child has not yet learned how to hold and empty urine well. (Communication between the brain and bladder may take time to develop.)
- Your child's body makes too much urine at night.
- Your child is constipated. Full bowels can put pressure on the bladder and lead to problems with holding and emptying urine well.
- Your child has a minor illness, is overly tired, or is responding to changes or stresses going on at home.
- There is a family history of bedwetting. Most children who wet the bed have at least one parent who had the same problem as a child.
- Your child's bladder is small or not developed enough to hold urine for a full night.
- Your child has an underlying medical problem.

What you can do

Most children wet their beds during toilet training. Even after they stay dry at night for a number of days or even weeks, they may start wetting at night again. If this happens to your child, simply go back to training pants at night and try again another time. The problem usually disappears as children get older. If children reach school age and still have

problems wetting the bed, it most likely means they have never developed nighttime bladder control.

Signs of a medical problem

If your child has been completely toilet trained for 6 months or longer and suddenly begins wetting the bed, talk with your child's provider. It may be a sign of a medical problem. However, most medical problems that cause bedwetting to recur suddenly have other signs, including:

- Changes in how much and how often your child urinates during the day
- Pain, burning, or straining while urinating
- A very small or narrow stream of urine or dribbling that is constant or happens just after urination
- Cloudy or pink urine or bloodstains on underpants
- Daytime and nighttime wetting
- Sudden change in personality or mood
- Poor bowel control
- Urinating after stress (coughing, running, or lifting)
- Certain gait disturbances (problems with walking that may mean an underlying neurologic problem)
- Continuous dampness

Managing bedwetting

Keep the following tips in mind when dealing with bedwetting:

- Do not blame your child. Remember that it is not your child's fault.
- Be honest with your child about what is going on. Let your child know it's not his or her fault and that most children outgrow bedwetting.
- Be sensitive to your child's feelings. If you don't make a big issue out of bedwetting, chances are your child won't either. Also remind your child that other children wet the bed.
- Protect the bed. A plastic cover under the sheets protects the mattress from getting wet and smelling like urine.
- Let your child help. Encourage your child to help change the wet sheets and covers. This teaches responsibility. It can also keep your child from feeling embarrassed if the rest of the family knows. However, if your child sees this as punishment, it is not recommended.
- Set a no-teasing rule in your family. Do not let family members, especially siblings, tease your child. Let them know that it's not your child's fault.
- Take steps before bedtime. Have your child use the toilet and avoid drinking large amounts of fluid just before bedtime.
- Try to wake your child up to use the toilet 1 to 2 hours after going to sleep to help your child stay dry through the night.
- Be positive. Reward your child for dry nights. Offer support, not punishment, for wet nights.
- Be aware of your child's daily urine and bowel habits.

Bedwetting alarms

If your child is still not able to stay dry during the night after using these steps for 1 to 3 months, a bedwetting alarm may be recommended. When a bedwetting alarm senses urine, it sets off an alarm so the child can wake up to use the toilet. When used correctly, it will detect wetness right away and sound the alarm. Be sure your child resets the alarm before going back to sleep.

Bedwetting alarms are successful 50% to 75% of the time. They tend to be most helpful for children who are deep sleepers and have some bladder control on their own. Ask your child's doctor which type of alarm would be best for your child.

Medicines

Medicines are available to treat bedwetting for children 6 years and older. Though medicines rarely cure bedwetting, they may be helpful, especially when children begin attending sleepovers or overnight camps. Your child's doctor can tell you more about these medicines and if they are right for your child. Remember to ask about possible side effects.

Source

<https://www.healthychildren.org/English/ages-stages/toddler/toilet-training/Pages/Bedwetting.aspx>